ELSEVIER

Gene 180 (1996) 229–231

GENE
AN INTERNATIONAL JOURNAL ON GENES AND GENOMES

Author Index

Agou, F., see Lazard, M. (180) 197
Ajdić, D., Sutcliffe, I.C., Russell, R.R.B. and Ferretti, J.J.
Organization and nucleotide sequence of the *Streptococcus mutans* galactose operon (180) 137
Alagón, A., see Miranda, R. (180) 37
Alborn, W.E., Jr., Hoskins, J.A., Ünal, S., Flokowitsch, J.E., Hayes, C.A., Dotzlaf, J.E., Yeh, W.K. and Skatrud, P.L.
Cloning and characterization of *femA* and *femB* from *Staphylococcus epidermidis* (180) 177
Aliyeva, E., Metz, A.M. and Browning, K.S.
Sequences of two expressed sequence tags (EST) from rice encoding different cap-binding proteins (180) 221
Alm, R.A., see Watson, A.A. (180) 49
Altenbuchner, J., see Fischer, J. (180) 81
Anderson, S.K., see Nestel, F.P. (180) 151
Aoshima, M., Nishibe, Y., Hasegawa, M., Yamagishi, A. and Oshima, T.
Cloning and sequencing of a gene encoding 16S ribosomal RNA from a novel hyperthermophilic archaebacterium NC12 (180) 183
Azakami, H., Yumoto, H., Nakae, H., Matsuo, T. and Ebisu, S.
Molecular analysis of the gene encoding a protein component of the *Eikenella corrodens* adhesin complex that is close to the carbohydrate recognition domain (180) 207

Beldjord, C., see Vázquez, M.P. (180) 43
Benito, A., see Villaverde, A. (180) 101
Benner, J.S., see Guthrie, E.P. (180) 107
Bidwell, C.A., see Reecy, J.M. (180) 23
Bienvenu, T., see Vázquez, M.P. (180) 43
Boucher, H., Mercure, S., Montplaisir, S. and Lemay, G.
A novel group I intron in *Candida dubliniensis* is homologous to a *Candida albicans* intron (180) 189
Bragonzi, A., see Monaco, L. (180) 145
Briley, G.P., see Reecy, J.M. (180) 23
Brooks, J.E., see Guthrie, E.P. (180) 107
Browning, K.S., see Aliyeva, E. (180) 221
Buchholz, M.A., see Kwon, T.K. (180) 113

Cavarelli, J., see Lazard, M. (180) 197
Chan, L., see Oka, K. (180) 69
Chen, P., see Rich, T. (180) 125
Chen, W., Yap, S.F. and Lim, L.
Isolation of the gene coding for *Caenorhabditis elegans* Rac2 homologue, a Ras-related small GTP-binding protein (180) 217
Chiu, I.-M., see Hackshaw, K.V. (180) 131
Chu, M.-j., see Oka, K. (180) 69
Colwill, K., see Nestel, F.P. (180) 151
Coote, J.G., see Westrop, G.D. (180) 91
Copeland, N.G., see Luo, C. (180) 29
Coutelle, C., see Villaverde, A. (180) 101
Cundliffe, E., see Gandecha, A.R. (180) 173

Da Costa, N.A., see Westrop, G.D. (180) 91
Disteche, C., see Luo, C. (180) 29
Dombrowski, S.M., see Maitra, S. (180) 165
Dotzlaf, J.E., see Alborn, W.E. Jr. (180) 177

Ebisu, S., see Azakami, H. (180) 207
Edelhoff, S., see Luo, C. (180) 29

Feliu, J.X., see Villaverde, A. (180) 101
Ferretti, J.J., see Ajdić, D. (180) 137
Fischer, J., Maier, H., Viell, P. and Altenbuchner, J.
The use of an improved transposon mutagenesis system for DNA sequencing leads to the characterization of a new insertion sequence of *Streptomyces lividans* 66 (180) 81
Flokowitsch, J.E., see Alborn, W.E. Jr. (180) 177
Frandsen, N., see Guérout-Fleury, A.-M. (180) 57
Furlow, Z.M., see Hackshaw, K.V. (180) 131
Furman, F., see Rich, T. (180) 125

Gandecha, A.R. and Cundliffe, E.
Molecular analysis of *tlrD*, an MLS resistance determinant from the tylosin producer, *Streptomyces fradiae* (180) 173
Ganguly, R., see Maitra, S. (180) 165
García, J.L., see Sánchez-Beato, A.R. (180) 13
Giovanazzi, S., see Monaco, L. (180) 145
Grant, A.L., see Reecy, J.M. (180) 23
Guérout-Fleury, A.-M., Frandsen, N. and Stragier, P.
Plasmids for ectopic integration in *Bacillus subtilis* (180) 57
Guthrie, E.P., Quinton-Jager, T., Moran, L.S., Slatko, B.E., Kucera, R.B., Benner, J.S., Wilson, G.G. and Brooks, J.E.
Cloning, expression and sequence analysis of the *Sph*I restriction-modification system (180) 107

Hackshaw, K.V., Furlow, Z.M. and Chiu, I.-M.
Cloning and characterization of a novel upstream untranslated exon of the mouse *Fgf*-1 gene (180) 131
Hamajima, N., Matsuda, K., Sakata, S., Tamaki, N., Sasaki, M. and Nonaka, M.
A novel gene family defined by human dihydropyrimidinase and three related proteins with differential tissue distribution (180) 157
Harbottle, R.P., see Villaverde, A. (180) 101
Harper, S., see Nestel, F.P. (180) 151
Hasegawa, M., see Aoshima, M. (180) 183
Hayes, C.A., see Alborn, W.E. Jr. (180) 177
Herzog, C.R., see Soloff, E.V. (180) 213
Hofmann, K.J., see Neeper, M.P. (180) 1
Hogan, P.G., see Luo, C. (180) 29
Hormozi, E.K., see Westrop, G.D. (180) 91
Hoskins, J.A., see Alborn, W.E. Jr. (180) 177
Huynh, N., see Rich, T. (180) 125

Ishihama, A., see Sakurai, H. (180) 63
Ishimura-Oka, K., see Oka, K. (180) 69
Israel, M.A., see Rich, T. (180) 125

Jansen, K.U., see Neeper, M.P. (180) 1
Jenkins, N.A., see Luo, C. (180) 29

Kilby, N.J., see Snaith, M.R. (180) 225
Krebber, H. and Ponstingl, H.
Ubiquitous expression and testis-specific alternative polyadenylation of mRNA for the human Ran GTPase activator RanGAP1 (180) 7
Kucera, R.B., see Guthrie, E.P. (180) 107
Kwon, T.K., Nagel, J.E., Buchholz, M.A. and Nordin, A.A.
Characterization of the murine cyclin-dependent kinase inhibitor gene *p27^{Kip1}* (180) 113

Latreille, M.-T., see Lazard, M. (180) 197
Lazard, M., Agou, F., Cavarelli, J., Latreille, M.-T., Moras, D. and Mirande, M.
Genomic organization of the rat aspartyl-tRNA synthetase gene family: a single active gene and several retropseudogenes (180) 197
Lemay, G., see Boucher, H. (180) 189
Levín, M.J., see Vázquez, M.P. (180) 43
Lijam, N., see Yang, Y. (180) 121
Lim, L., see Chen, W. (180) 217
Lizardi, P.M., see Miranda, R. (180) 37
Lorenzi, H., see Vázquez, M.P. (180) 43
Luo, C., Copeland, N.G., Jenkins, N.A., Edelhoff, S., Disteche, C., Hogan, P.G. and Rao, A.
Normal function of the transcription factor NFAT1 in *wasted* mice. Chromosome localization of NFAT1 gene (180) 29

Maier, H., see Fischer, J. (180) 81
Maitra, S., Dombrowski, S.M., Waters, L.C. and Ganguly, R.
Three second chromosome-linked clustered *Cyp6* genes show differential constitutive and barbital-induced expression in DDT-resistant and susceptible strains of *Drosophila melanogaster* (180) 165
Matsuda, K., see Hamajima, N. (180) 157
Matsuo, T., see Azakami, H. (180) 207
Mattick, J.S., see Watson, A.A. (180) 49
Mercure, S., see Boucher, H. (180) 189
Metz, A.M., see Aliyeva, E. (180) 221
Miranda, R., Salgado, L.M., Sánchez-López, R., Alagón, A. and Lizardi, P.M.
Identification and analysis of the *u6 small nuclear RNA* gene from *Entamoeba histolytica* (180) 37
Mirande, M., see Lazard, M. (180) 197
Miyao, T., see Sakurai, H. (180) 63
Monaco, L., Tagliabue, R., Giovanazzi, S., Bragonzi, A. and Soria, M.R.
Expression of recombinant human granulocyte colony-stimulating factor in CHO dhfr$^-$ cells: new insights into the in vitro amplification expression system (180) 145
Montplaisir, S., see Boucher, H. (180) 189
Moran, L.S., see Guthrie, E.P. (180) 107
Moras, D., see Lazard, M. (180) 197
Murray, J.A.H., see Snaith, M.R. (180) 225

Nagel, J.E., see Kwon, T.K. (180) 113
Nakae, H., see Azakami, H. (180) 207
Neeper, M.P., Hofmann, K.J. and Jansen, K.U.
Expression of the major capsid protein of human papillomavirus type 11 in *Saccharomyces cerevisae* (180) 1
Nestel, F.P., Colwill, K., Harper, S., Pawson, T. and Anderson, S.K.
RS cyclophilins: Identification of an NK-TR$_1$-related cyclophilin (180) 151
Nishibe, Y., see Aoshima, M. (180) 183
Nonaka, M., see Hamajima, N. (180) 157
Nordin, A.A., see Kwon, T.K. (180) 113

Oka, K., Ishimura-Oka, K., Chu, M.-j. and Chan, L.
Transcription of the human hepatic lipase gene is modulated by multiple negative elements in HepG2 cells (180) 69
Oshima, T., see Aoshima, M. (180) 183

Parton, R., see Westrop, G.D. (180) 91
Pawson, T., see Nestel, F.P. (180) 151
Ponstingl, H., see Krebber, H. (180) 7

Quinton-Jager, T., see Guthrie, E.P. (180) 107

Rao, A., see Luo, C. (180) 29
Reecy, J.M., Bidwell, C.A., Briley, G.P. and Grant, A.L.
Structure and regulation of the porcine skeletal α-actin-encoding gene (180) 23
Rich, T., Chen, P., Furman, F., Huynh, N. and Israel, M.A.
RTVP-1, a novel human gene with sequence similarity to genes of diverse species, is expressed in tumor cell lines of glial but not neuronal origin (180) 125
Russell, R.R.B., see Ajdić, D. (180) 137

Sakata, S., see Hamajima, N. (180) 157
Sakurai, H., Miyao, T. and Ishihama, A.
Subunit composition of RNA polymerase II from the fission yeast *Schizosaccharomyces pombe* (180) 63
Salgado, L.M., see Miranda, R. (180) 37
Sánchez-Beato, A.R. and García, J.L.
Molecular characterization of a family of choline-binding proteins of *Clostridium beijerinckii* NCIB 8052. Evolution and gene redundancy in prokaryotic cell (180) 13
Sánchez-López, R., see Miranda, R. (180) 37
Sasaki, M., see Hamajima, N. (180) 157
Skatrud, P.L., see Alborn, W.E. Jr. (180) 177
Slatko, B.E., see Guthrie, E.P. (180) 107
Snaith, M.R., Kilby, N.J. and Murray, J.A.H.
An *Escherichia coli* system for assay of Flp site-specific recombination on substrate plasmids (180) 225
Soloff, E.V., Herzog, C.R. and You, M.
The 5′-flanking region of the E1 α form of the murine p16^{INK4a} (MTS1) gene (180) 213
Soria, M.R., see Monaco, L. (180) 145
Stragier, P., see Guérout-Fleury, A.-M. (180) 57
Sussman, D.J., see Yang, Y. (180) 121
Sutcliffe, I.C., see Ajdić, D. (180) 137

Tagliabue, R., see Monaco, L. (180) 145
Tamaki, N., see Hamajima, N. (180) 157
Tsang, M., see Yang, Y. (180) 121

Ünal, S., see Alborn, W.E. Jr. (180) 177

Vázquez, M.P., Beldjord, C., Lorenzi, H., Bienvenu, T. and Levín, M.J.
Detection of polymorphism in the *Trypanosoma cruzi* TcP2β gene family by single strand conformational analysis (SSCA) (180) 43
Viell, P., see Fischer, J. (180) 81
Villaverde, A., Feliu, J.X., Harbottle, R.P., Benito, A. and Coutelle, C.
A recombinant, arginine-glycine-aspartic acid (RGD) motif from foot-and-mouth disease virus binds mammalian cells through vitronectin and, to a lower extent, fibronectin receptors (180) 101

Waters, L.C., see Maitra, S. (180) 165
Watson, A.A., Alm, R.A. and Mattick, J.S.
Identification of a gene, *pilF*, required for type 4 fimbrial biogenesis and twitching motility in *Pseudomonas aeruginosa* (180) 49
Westrop, G.D., Hormozi, E.K., Da Costa, N.A., Parton, R. and Coote, J.G.
Bordetella pertussis adenylate cyclase toxin: proCyaA and CyaC

proteins synthesised separately in *Escherichia coli* produce active toxin in vitro (180) 91
Wilson, G.G., see Guthrie, E.P. (180) 107

Yamagishi, A., see Aoshima, M. (180) 183
Yang, Y., Lijam, N., Sussman, D.J. and Tsang, M.
Genomic organization of mouse *Dishevelled* genes (180) 121
Yap, S.F., see Chen, W. (180) 217
Yeh, W.K., see Alborn, W.E. Jr. (180) 177
You, M., see Soloff, E.V. (180) 213
Yumoto, H., see Azakami, H. (180) 207